

Product Information

Lectin
from *Bandeiraea simplicifolia*
BS-I Isolectin B4
Biotin Conjugate

Catalog Number **L2140**
Storage Temperature –20 °C

Product Description

Lectins are proteins or glycoproteins of non-immune origin that agglutinate cells and/or precipitate complex carbohydrates. Lectins are capable of binding glycoproteins even in presence of various detergents.¹ The agglutination activity of these highly specific carbohydrate-binding molecules is usually inhibited by a simple monosaccharide, but for some lectins, di, tri, and even polysaccharides are required.

Lectins are isolated from a wide variety of natural sources, including seeds, plant roots and bark, fungi, bacteria, seaweed and sponges, mollusks, fish eggs, body fluids of invertebrates and lower vertebrates, and from mammalian cell membranes. The precise physiological role of lectins in nature is still unknown, but they have proved to be very valuable in a wide variety of applications *in vitro*, including:

1. blood grouping and erythrocyte polyagglutination studies.
2. mitogenic stimulation of lymphocytes.
3. lymphocyte subpopulation studies.
4. fractionation of cells and other particles.
5. histochemical studies of normal and pathological conditions.

Sigma offers a range of lectins suitable for the above applications. Most Sigma lectins are highly purified by affinity chromatography, but some are offered as purified or partially purified lectins, suitable for specific applications.

Many of the lectins are available conjugated to (conjugation does not alter the specificity of the lectin):

1. fluorochromes (for detection by fluorimetry).
2. enzymes (for enzyme-linked assays).
3. insoluble matrices (for use as affinity media).

Please refer to the table for general information on the most common lectins.

This product is conjugated to biotin, allowing detection in binding assays through a secondary reporter such as streptavidin-peroxidase.

Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Preparation Instructions

The product is soluble in 0.9% sodium chloride solution (1 mg/ml), yielding a clear, yellow solution.

Storage/Stability

Aggregation is thought to occur in the presence of high concentrations of 2-mercaptoethanol.

| Lectin | MW (kDa) | Subunits | Specificity | | Mitogenic Activity |
|-----------------------------------|----------------------|-----------------|--------------------|-----------------------|---------------------------|
| | | | Blood Group | Sugar | |
| <i>Abrus precatorius</i> | | | — | gal | + |
| Agglutinin | 134 | 4 | | gal | |
| Abrin A (toxin) | 60 | 2 | | gal | |
| Abrin B (toxin) | 63.8 | 2(αβ) | | gal | |
| <i>Agarius bisporus</i> | 58.5 | — | — | β-gal(1→3)galNAc | |
| <i>Anguilla anguilla</i> | 40 | 2 | H | α-L-Fuc | |
| <i>Arachis hypogaea</i> | 120 | 4 | T | β-gal(1→3)galNAc | |
| <i>Artocarpus integrifolia</i> | 42 | 4 | T | α-gal→OMe | + |
| <i>Bandeiraea simplicifolia</i> | | | | | |
| BS-I | 114 | 4 | A, B | α-gal, α-galNAc | |
| BS-I-A ₄ | 114 | 4 | A | α-galNAc | |
| BS-I-B ₄ | 114 | 4 | B | α-gal | |
| BS-II | 113 | 4 | acq, B, Tk, T | glcNAc | |
| <i>Bauhinia purpurea</i> | 195 | 4 | — | β-gal(1→3)galNAc | + |
| <i>Caragana arborescens</i> | 60; 120 ^a | 2/4 | — | galNAc | |
| <i>Cicer arietinum</i> | 44 | 2 | — | fetuin | |
| <i>Codium fragile</i> | 60 | 4 | — | galNAc | |
| <i>Concanavalin A</i> | 102 | 4 | — | α-man, α-glc | + |
| <i>Succinyl-Concanavalin A</i> | 51 | 2 | — | α-man, α-glc | + |
| <i>Cytisus scoparius</i> | — | — | — | galNAc, gal | |
| <i>Datura stramonium</i> | 86 | 2(αβ) | — | (glcNAc) ₂ | |
| <i>Dolichos biflorus</i> | 140 | 4 | A ₁ | α-galNAc | |
| <i>Erythrina corallodendron</i> | 60 | 2 | — | β-gal(1→4)glcNAc | + |
| <i>Erythrina cristagalli</i> | 56.8 | 2(αβ) | — | β-gal(1→4)glcNAc | |
| <i>Euonymus europaeus</i> | 166 | 4(αβ) | B, H | α-gal(1→3)gal | + |
| <i>Galanthus nivalis</i> | 52 | 4 | (h) | non-reduc. α-man | |
| <i>Glycine max</i> | 110 | 4 | — | galNAc | + |
| <i>Helix aspersa</i> | 79 | — | A | galNAc | |
| <i>Helix pomatia</i> | 79 | 6 | A | galNAc | |
| <i>Lathyrus odoratus</i> | 40-43 | 4(αβ) | — | α-man | + |
| <i>Lens culinaris</i> | 49 | 2 | — | α-man | + |
| <i>Limulus polyphemus</i> | 400 | 18 | — | NeuNAc | |
| Bacterial agglutinin | — | — | — | galNAc, glcNAc | |
| <i>Lycopersicon esculentum</i> | 71 | — | — | (glcNAc) ₃ | |
| <i>Maackia amurensis</i> | 130 | 2(αβ) | O | sialic acid | + |
| <i>Maclura pomifera</i> | 40-43 | 2(αβ) | — | α-gal, α-galNAc | |
| <i>Momordica charantia</i> | 115-129 | 4(αβ) | — | gal, galNAc | |
| <i>Naja mocambique mocambique</i> | — | — | — | — | |
| <i>Naja naja kaouthia</i> | — | — | — | — | |
| <i>Narcissus pseudonarcissus</i> | 26 | 2 | (h) | α-D-man | |
| <i>Persea americana</i> | — | — | — | — | |
| <i>Phaseolus coccineus</i> | 112 | 4 | — | — | |
| <i>Phaseolus limensis</i> | 247(II) | 8 | A | galNAc | + |
| | 124(III) | 4 | | | |
| <i>Phaseolus vulgaris</i> | | | | | |
| PHA-E | 128 | 4 | — | oligosaccharide | + |
| PHA-L | 128 | 4 | — | oligosaccharide | + |
| PHA-P | | | | | |
| PHA-M | | | | | |

----- Table continued on next page -----

| Lectin | MW (kDa) | Subunits | Specificity | | Mitogenic Activity |
|------------------------------------|----------------------|-----------------|--------------------------------|--------------------------------|---------------------------|
| | | | Blood Group | Sugar | |
| <i>Phytolacca americana</i> | 32 | — | — | (glcNAc) ₃ | + |
| <i>Pisum sativum</i> | 49 | 4(αβ) | — | α-man | + |
| <i>Pseudomonas aeruginosa PA-I</i> | 13-13.7 | — | — | gal | + ^c |
| <i>Psophocarpus tetragonolobus</i> | 35 | 1 | — | galNAc, gal | |
| <i>Ptilota plumosa</i> | 65; 170 | — | B | α-gal | |
| <i>Ricinus communis</i> | | | | | |
| Toxin, RCA ₆₀ | 60 | 2 | — | galNAc, β-gal | |
| Toxin, RCA ₁₂₀ | 120 | 4 | — | β-gal | |
| <i>Sambucus nigra</i> | 140 | 4(αβ) | — | αNeuNAC(2→6)gal | + ^c |
| | | | | galNAc | |
| <i>Solanum tuberosum</i> | 50; 100 ^a | 1, 2 | — | (glcNAc) ₃ | |
| <i>Sophora japonica</i> | 133 | 4 | A, B | β-galNAc | |
| <i>Tetragonolobus purpureas</i> | 120(A) | 4 | H | α-L-fuc | |
| | 58(BA) | 2 | H | α-L-fuc | |
| | 117(C) | 4 | H | α-L-fuc | |
| <i>Triticum vulgaris</i> | 36 | 2 | — | (glcNAc) ₂ , NeuNAc | + |
| <i>Ulex europaeus</i> | | | | | |
| UEA I | 68 | — | H | α-L-fuc | |
| UEA II | 68 | — | — | (glcNAc) ₂ | |
| <i>Vicia faba</i> | 50 | 4(αβ) | — | man, glc | + |
| <i>Vicia sativa</i> | 40 | 4(αβ) | — | glc, man | + |
| <i>Vicia villosa</i> | 139 | 4 | A ₁ +T _n | galNAc | |
| A ₄ | 134 | 4 | A ₁ | galNAc | |
| B ₄ | 143 | 4 | T _n | galNAc | |
| <i>Vigna radiata</i> | 160 | 4 | — | α-gal | |
| <i>Viscum album</i> | 115 | 4(αβ) | — | β-gal | |
| <i>Wisteria floribunda</i> | 68 | 2 | — | galNAc | |

^a Concentration-dependent molecular weight

^b Non-agglutinating and mitogenic

^c Mitogenic for neuraminidase-treated lymphocytes

References

1. Reisfeld, R. A., et al., Isolation and Characterization of a Mitogen from Pokeweed (*Phytolacca americana*). Proc. Nat. Acad. Sci., **58(5)**, 2020-2027 (1967).
2. Mansour, H. et al., Connexin 30 expression and frequency of connexin heterogeneity in astrocyte gap junction plaques increase with age in the rat retina. PLoS ONE, 2013 Mar 14, 8(3), e57038.

IRB,MWM,JRC,NSB,SAG,AI,MAM 03/16-1