

3050 Spruce Street, St. Louis, MO 63103 USA
Tel: (800) 521-8956 (314) 771-5765 Fax: (800) 325-5052 (314) 771-5757
email: techservice@sial.com sigma-aldrich.com

Product Information

Cell freezing Media

sterile-filtered, suitable for cell culture

Product Number **C6039**Cell Freezing Medium-Glycerol 1×
Product Number **C6164**Cell Freezing Medium-DMSO 1×
Product Number **C6295**Cell Freezing Medium-DMSO Serum free 1×
Storage Temperature –20 °C

Product Description

The Cell Freezing Media are complete ready-to-use reagents which are designed to protect and preserve cells during frozen storage.

Product Numbers C6039 and C6164 contain Minimum Essential Medium (MEM) supplemented with a mixture of fetal bovine serum and calf serum, and contain 10% glycerol or DMSO.

Product Number C6295 is a serum-free preparation prepared according to the original published formula of Charity Waymouth.¹ It contains 8.7% DMSO in MEM supplemented with methyl cellulose.

Precautions and Disclaimer

For R&D use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

Cell Freezing Media should be stored at -20 °C. After thawing, these products may be stored at 2-8 °C for up to 5 days. It is not recommend to store reagents in a frost-free freezer as temperature cycling may cause deterioration of the products.

Procedure

Cell Freezing Media may be used in standard freezing protocols. The following protocol may be used:

- Thaw the Cell Freezing Medium and hold it on wet ice.
- Remove adherent cells with trypsin or other appropriate means. For optimal results cells should be in the log phase of growth.

- 3. Gently pellet (10 minutes at $250 \times g$ at 2–8 °C) the cells by centrifugation and remove as much of the growth medium as possible.
- Suspend the cells in freezing medium at 10⁶ to 10⁷ cells/ml. Myelomas or hybridomas may require a higher cell density.
- 5. Aliquot cells into freezing vials, holding them on wet ice until freezing begins (within 5 minutes).
- Freeze cells according to standard protocols. Store below –70 °C.

Thawing of frozen cells may be done as follows:

- 1. Remove cells from frozen storage and quickly thaw in a 37 °C water bath.
- 2. Dilute 1 ml of cell suspension with 10 ml of complete growth medium.
- 3. Gently mix and pellet cells by gentle centrifugation.
- 4. Suspend cells in complete growth medium and plate in appropriate vessels.

References

- Waymouth, C., and Varnum, D., (1976). Simple Freezing Procedure for Storage in Serum-Free Media of Cultured and Tumor Cells of Mouse. TCA Manual 2:1 pp. 311-313.
- Freshney, R., (1986). Animal Cell Culture: A Practical Approach. IRL Press Ltd., Oxford. pp. 73-77.
- 3. Kruse, P. and Patterson, M. (1973). Tissue Culture Methods and Applications. Academic Press, NY. pp. 712-718.

PC,MAM 06/19-1