

Product Information

Mucin from porcine stomach

Type III, bound sialic acid 0.5-1.5%, partially purified powder

M1778

Product Description

CAS Registry Number: 84062-64-4

Mucins are highly glycosylated proteins that occur in mucus and the mucous membranes of animals and humans. The principal glycoprotein component of mucus, mucins are associated with such organs as the abdomen and the stomach. Their high degree of glycosylation renders mucins highly stable to digestion in acidic *in vivo* environments such as gastric mucosa.

Polymeric mucin has been estimated to have a molecular mass of ~4,000-5,500 kDa.^{1,2} The molecular mass of mucin monomer has been estimated at ~640 kDa.² Estimates of porcine mucin composition include a carbohydrate portion of 80-84%, and a protein portion of 16-20%.¹⁻³ The peptide backbone of mucin is notably rich in Pro, Ser, and Thr residues, which comprise ~43% of the amino acid composition of the mucin peptide backbone. Cys residues are prevalent in the non-glycosylated portions of mucin, and allow for crosslinking of mucin monomers to create highly complex polymers.

In porcine mucin, *O*-glycosylation occurs particularly at serine and threonine residues. The more prevalent *O*-linked glycans in porcine mucin include:³

- N-acetylgalactosamine
- N-acetylglucosamine
- Fucose
- Galactose
- *N*-acetylneuraminic acid (sialic acid)

Several studies have investigated the structural properties of porcine gastric mucin, and proposed various structural models.⁴⁻⁶ Several theses⁷⁻⁹ and dissertations¹⁰⁻¹⁶ have cited use of product M1778 in their protocols.

Reagent

This product is a partially purified powder with bound sialic acid content in the range of 0.5-1.5%, measured as *N*-acetylneuraminic acid. This product is not sterile.

Precautions and Disclaimer

For R&D use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Procedure

A general procedure for sterilizing mucin has been published,¹⁷ as follows:

- 1. Place dry powdered mucin in a flask.
- 2. Cover the mucin with 95% ethanol.
- 3. Heat the covered mucin at 70 °C for 24 hours.
- 4. Evaporate the alcohol to recover the mucin.

However, we have not tested this procedure ourselves.

Storage/Stability

Store the product at 2-8 °C.

References

1

- Fogg, F.J.J. et al., Biochem. J., 316(Pt 3), 937-942 (1996).
- 2. Schömig, V.J. *et al.*, *RSC Adv.*, **6**, 44932-44943 (2016).
- 3. Libao-Mercado, A.J., and de Lange, C.F.M., *Livestock Sci.*, **109(1-3)**, 141-144 (2007).



- Perez-Vilar, J., and Hill, R.L., J. Biol. Chem., 274(35), 31751-31754 (1999).
- Di Cola, E. et al., Biomacromolecules, 9(11), 3216-3222 (2008).
- Veluraja, K. et al., Biochem. Biophys. Res. Comm., 406(94), 570-573 (2011).
- 7. Kore, Doris Rachel, "In-Vitro Comparison of Tensile Bond Strength of Denture Adhesives on Denture Bases". Loma Linda University, M.S. thesis, p. 12 (2012).
- Owuor, Daniel R., "A Study of Cytocompatibility and Biodegradability of Az31 Magnesium Alloy in Simulated Airway Surface Lining Fluid". North Carolina Agricultural and Technical State University, M.S. thesis, p. 27 (2013).
- 9. Tam, Ngai Keung, "Comfort and Compatibility of Silicone Hydrogel Contact Lenses". Brigham Young University, M.S. thesis, p. 16 (2013).
- 10. Jumel, Kornelia, "Molecular Size of Interacting and Degrading Polysaccharides". University of Nottingham, Ph.D. dissertation, p. 102 (1994).
- 11. Sandberg, Tomas, "On the Development of Mucin-based Biomaterial Coatings". Uppsala Universitet, Ph.D. dissertation, p. 31 (2008).
- 12. Kraatz, Mareike, "Isolation of lactic acid-related bacteria from the pig mucosal proximal gastrointestinal tract, including Olsenella umbonata sp. nov. and Veillonella magna sp. nov.". Freien Universität Berlin, Dr. vet. med. dissertation, pp. 79, 80 (2011).
- 13. Woo, Myung-Heui, "Improving protection against viral aerosols through development of novel decontamination methods and characterization of viral aerosol". University of Florida, Ph.D. dissertation, pp. 27, 106 (2012).
- 14. Robinson, Camilla, "Utilisation of mucin sulphur by *Pseudomonas aeruginosa* Importance for cystic fibrosis". University of Manchester, Ph.D. dissertation, p. 52 (2013).

- 15. Ferreira, William Thomas, "Clostridium difficile infection; establishment, prevention and the role of the microbiota". Royal Holloway, University of London, Ph.D. dissertation, p. 71 (2019).
- 16. Marczynski, Matthias, "Structure-function relationships of mucin glycoproteins". Technische Universität München, Dr. rer. nat. dissertation, pp. 9, 64 (2022).
- 17. Yeung, A.T.Y. *et al.*, *MBio*, **3(3)**, e00073-12 (2012).

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at SigmaAldrich.com/techservice.

Standard Warranty

The applicable warranty for the products listed in this publication may be found at SigmaAldrich.com/terms.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

The life science business of Merck operates as MilliporeSigma in the U.S. and Canada.

Merck and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

