

3050 Spruce Street Saint Louis, Missouri 63103 USA Telephone 800-325-5832 • (314) 771-5765 Fax (314) 286-7828 email: techserv@sial.com sigma-aldrich.com

# ProductInformation

Pyruvate Kinase preparation from rabbit muscle

Product Number **P 9136** Storage Temperature -0 °C

## **Product Description**

CAS Number: 9001-59-6 Enzyme Commission (EC) Number: 2.7.1.40 Molecular Weight: 237 kDa<sup>1</sup> Extinction Coefficient:  $E^{0.1\%} = 0.54^{1}$ 

Pyruvate kinase from rabbit muscle is a tetramer consisting of four equal subunits of molecular weight 57 kDa.<sup>1</sup>

Pyruvate kinase catalyzes the following reaction:

ATP + Pyruvate  $\rightarrow$  ADP + Phophoenolpuruvate

Reported K<sub>m</sub> values are ATP (0.86 mM), pyruvate (10 mM), ADP (0.3 mM), and PEP (0.07 mM).<sup>2</sup> Pyruvate kinase can also utilize other dinucleotide phosphates as substrates including GDP, IDP, dADP, UDP, CDP, dCDP.<sup>1</sup> The pH optimum for pyruvate kinase is  $7.5^3$ , and the mechanism of the reaction catalyzed by pyruvate kinase has been described.<sup>4</sup>

Both  $Mg^{2+}$  and  $K^+$  are required metal cofactors for optimal activity. The enzyme is inhibited by  $Ca^{2+,1}$ .

Pyruvate kinase is often used as a coupling enzyme in conjunction with lactic dehydrogenase in quantifying ADP and the activity of enzymes that catalyze the formation of ADP.

## **Precautions and Disclaimer**

For Laboratory Use Only. Not for drug, household or other uses.

### **Preparation Instructions**

This enzyme is soluble in water (1 mg/ml), yielding a clear and colorless solution.

### References

- Kayne, F. J., in The Enzymes, 3rd ed., Vol. 8, pt. A, Boyer, P. D., ed., Academic press (New York, NY: 1973), pp. 353-382.
- Methods of Enzymatic Analysis, 2nd English ed., Vol. 1, Bergmeyer, H. U., ed., Academic Press (New York, NY: 1974), pp. 509-511.
- 3. Bucher, T., and Pfleiderer, G., Pyruvate kinase from muscle. Meth. Enzymol., 1, 435-440 (1955).
- Ainsworth, S., et al., The regulatory properties of rabbit muscle pyruvate kinase. The influence of substrate concentrations. Biochem. J., 209(2), 401-411 (1983).

TMG/RXR 1/04

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.