

Product Information

Guanidine hydrochloride solution, 6M

SRE0066

Product Description

This product is a ready-to-use 6 M guanidine hydrochloride solution. Guanidine hydrochloride is commonly used as a denaturant, because of its ability to break hydrogen bonds between amino acid residues. By breaking these bonds, the 3D conformation of the protein is unfolded and the aqueous solubility of the protein is greatly increased. Once denatured, the protein can be easily reduced, modified, or analyzed, in a variety of procedures. Further processing of the sample can allow the proteins to be analyzed by other common protein analysis methods including mass spectrometry, electrophoresis, and enzymatic digests. Guanidine hydrochloride has also typically been used for the isolation of RNA, to denature globular proteins, and for protein refolding studies. It can also be used to facilitate the generation of tryptic peptides for analysis of complex protein samples.

Precautions and Disclaimer

For manufacturing, processing, or repacking. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Preparation Instructions

The product is a clear, colorless liquid and is supplied ready-to-use. The 6 M Guanidine hydrochloride solution may be used as a stock solution to prepare solutions of lower concentrations as appropriate (see Table 1).

Note: If precipitation is present in the 6 M buffer, warm the bottle to 37 °C and mix until completely dissolved prior to use.

Table 1. Dilution Table - Dilutions are based on a 10 mL volume of the 6 M Guanidine HCl solution.

Desired Molarity	Volume of water to add to 10 mL of 6 M Guanidine HCl solution	Final Volume
6 M	0.0 mL	10.0 mL
5 M	2.0 mL	12.0 mL
4 M	4.0 mL	14.0 mL
3 M	10.0 mL	20.0 mL
2 M	20.0 mL	30.0 mL
1 M	50.0 mL	60.0 mL

Storage/Stability

Store at Room Temperature

The product is stable for two years from the date of manufacture when stored at room temperature. Do not use past expiration date printed on product label.

References

1. Bruice, P.Y., Organic Chemistry, 3rd Ed., Prentice Hall (Upper Saddle River, New Jersey: 2001), pp 950-953.
2. Cox, R.A., The Use of Guanidinium Chloride in the Isolation of Nucleic Acids, Methods in Enzymology, 12 B, 120-129 (1968).
3. Voet, D. et al., Fundamentals of Biochemistry, Upgrade Edition, John Wiley & Sons Inc (New York, New York: 2002), pp 151-152.
4. Lodish et al., Molecular Cell Biology, 4th Ed., W.H. Freeman and Company, (New York, New York: 2000). p 63.
5. Gygi, S.P. et al., Quantitative Analysis of Complex Protein Mixtures Using Isotope-coded Affinity Tags. Nature Biotechnology, 17, 994-999 (1999).
6. West, S.M. et al., Improved Protein Refolding Using Hollow-Fibre Membrane Dialysis. Biotechnol. Bioeng., 57, 590-599 (1998).
7. Lilie, H. et al., Advances in Refolding of Proteins Produced in E. Coli. Current Opinion in Biotechnology, 9, 497-501 (1998).

-
8. Marston, F.A.O., and Hartley, D.L., Solubilization of Protein Aggregates. *Methods Enzymol.*, 182, 264- 276 (1990).

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at SigmaAldrich.com/techservice.

Terms and Conditions of Sale

Warranty, use restrictions, and other conditions of sale may be found at SigmaAldrich.com/terms.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

The life science business of Merck operates as MilliporeSigma in the U.S. and Canada.

Merck and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2023 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

SRE0066 Rev 12/23

