

Product Information

Chitinase from *Streptomyces griseus*

Free of DNA contaminants, suitable for Microbiome research

SAE0158

Product Description

E.C. 3.2.1.14

CAS RN 9001-06-3

Synonyms: Chitodextrinase, Poly(β -(1 \rightarrow 4)-[2-acetamido-2-deoxy-D-glucoside]) glycanohydrolase

Chitinase is an extracellular complex of enzymes that degrade chitin.¹ Chitinases have been detected in many microorganisms and in plants.² In fungi, chitinases assist in morphogenesis, to break down the inherent chitin content of fungal cell walls. Plant chitinases help in resistance to fungal attack and counteracting fungal growth, by targeting those same fungal cell walls. In bacteria, bacterial chitinases assist in utilizing chitin as a carbon source and as an energy source.^{2,3} *Streptomyces griseus* produces multiple chitinases of different molecular masses after growth induction with chitin as the carbon source.³

The enzymatic hydrolysis of chitin to *N*-acetyl-D-glucosamine involves two consecutive enzyme reactions:

- The first reaction, chitodextrinase-chitinase, is a poly(β -(1 \rightarrow 4)-[2-acetamido-2-deoxy-D-glucoside])-glycanohydrolase, which removes chitobiose units from chitin.
- The second activity is *N*-acetyl-glucosaminidase-chitobiase, which cleaves the disaccharide to its monomer subunits, *N*-acetyl-D-glucosamine.

The study of microbial communities has recently been revolutionized by the widespread adoption of culture-independent analytical techniques such as 16S rRNA gene sequencing and metagenomics. Since DNA contamination during sample preparation is a concern of these sequence-based approaches,⁴ DNA extraction reagents free of DNA contaminants are essential.

Purified Chitinase SAE0158 undergoes strict quality control testing to ensure the absence of detectable levels of contaminating DNA, using 35 cycles of PCR amplification of 16S and 18S rDNA, using universal primer sets.

The optimal reaction temperature is 37 °C.

Specific Activity: ≥ 1000 units/g protein

Unit Definition

Unit Definition: One unit will liberate 1.0 mg of *N*-acetyl-D-glucosamine from chitin per hour at pH 6.0 at 25 °C in a 2-hour assay.

Storage/Stability

This product, as supplied, should be stored at -20 °C.

Preparation Instructions

Solutions of chitinase can be prepared in DNA-free water. Stock solutions of chitinase can be stored at -20 °C in frozen aliquots.

Precautions and Disclaimer

For R&D use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

References

1. *CRC Handbook of Biochemistry and Molecular Biology: Proteins*, Vol. II, CRC Press, p. 405 (1976).
2. Kezuka, Y. *et al.*, *J. Mol. Biol.*, **358**(2), 472-484 (2006).
3. Tanabe, T. *et al.*, *J. Biosci. Bioeng.*, **89**(1), 27-32 (2000).
4. Eisenhofer, R. *et al.*, *Trends Microbiol.*, **27**(2), 105-117 (2019).

Notice

MERCK

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at SigmaAldrich.com/techservice.

Standard Warranty

The applicable warranty for the products listed in this publication may be found at SigmaAldrich.com/terms.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

The life science business of Merck operates
as MilliporeSigma in the U.S. and Canada.

Merck and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2021 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.
SAE0158pis Rev 10/21 EO,DT,GCY

