

3050 Spruce Street
Saint Louis, Missouri 63103 USA
Telephone 800-325-5832 • (314) 771-5765
Fax (314) 286-7828
email: techserv@sial.com
sigma-aldrich.com

# **ProductInformation**

## Actin from rabbit muscle

Product Number A 2522 Storage Temperature 2-8 °C

#### **Product Description**

CAS Number: 51005-14-2

Extinction coefficient:  $E^{1\%} = 1.11$  (G-actin),

1.17 (F-actin)

pl: 5.47 ( $\alpha$ ), 5.53 ( $\beta$ ), 5.50 ( $\gamma$ )<sup>6, 7, 8, 9</sup>

This material is prepared from skeletal muscle by using a modification of a reported method. <sup>1</sup> The molecular weight of the polypeptide by sequence analysis is 41.785 kDa and when calcium and ATP are included, the total is 42.3 kDa. <sup>2,3</sup> Methods have been reported for converting this product to the F-actin form. <sup>5</sup>

Monomeric rabbit actin has been used to inhibit DNAse I, with a binding constant =  $5 \times 10^8 \, \text{M}^{-1.10}$  The atomic structure of the actin:DNase I complex along with the amino acid sequence has been reported. It is noted that the ability of actin to polymerize is lost after forming a 1:1 complex with DNase I.

### **Precautions and Disclaimer**

For Laboratory Use Only. Not for drug, household or other uses.

#### **Preparation Instructions**

This material is soluble in water (1 mg/ml), yielding a hazy solution.

#### References

- Spudich, J. A., and Watt, S., The regulation of rabbit skeletal muscle contraction. I. Biochemical studies of the interaction of the tropomyosintroponin complex with actin and the proteolytic fragments of myosin. J. Biol. Chem., 246, 4866 (1971).
- 2. Elzinga, M., et al., Complete amino-acid sequence of actin of rabbit skeletal muscle. Proc. Natl. Acad. Sci. USA, **70**, 2687-2691 (1973).
- Collins, J.H., and Elzinga, M., The primary structure of actin from rabbit skeletal muscle. Completion and analysis of the amino acid sequence. J. Biol. Chem., 250, 5915-5920 (1975).
- 4. Houk, T. W. Jr., and Ue, K., The measurement of actin concentration in solution: a comparison of methods. Anal. Biochem., **62**, 66-74 (1974).
- 5. J. Chrom., **22**, 115 (1981).
- Zechel, K., and Weber, K., Actins from mammals, bird, fish and slime mold characterized by isoelectric focusing in polyacrylamide gels. Eur. J. Biochem., 89, 105-112 (1978).
- 7. Z. Physiol. Chem., 358, 1304 (1977).
- 8. Pardee, J.D., and Bamburg, J.R., Actin from embryonic chick brain. Isolation in high yield and comparison of biochemical properties with chicken muscle actin. Biochemistry, **18**, 2245 (1979).
- Flanagan, M.D., and Lin, S., Comparative studies on the characteristic properties of two forms of brain actin separable by isoelectric focusing.
   J. Neurochem., 32, 1037-1046 (1979).
- Mannherz, H.G., et al., The interaction of bovine pancreatic deoxyribonuclease I and skeletal muscle actin. Eur. J. Biochem., 104, 367-379 (1980).
- 11. Kabsch, W., et al., Atomic structure of the actin:DNase I complex. Nature, **347**, 37-44 (1990).

AGW/AJH 5/07